

UR-1537
ER/40685/919
July 1998

Gauge Mediated Supersymmetry Breaking and Spontaneous CP Violation as a Solution to the Strong CP Problem*

Otto C.W. Kong

*Department of Physics and Astronomy,
University of Rochester, Rochester NY 14627-0171.*

Abstract

The origin of CP violation is a major mystery, especially in relation to the strong CP problem. CP being a spontaneously broken symmetry could provide an elegant solution. However, such models have difficulty making themselves compatible with low-energy supersymmetry, which is popularly accepted as the solution to the hierarchy problem. We demonstrate that a certain class of low scale supersymmetric “Nelson-Barr” type models can solve the strong and supersymmetric CP problems while at the same time generating sufficient weak CP violation in the $K^0 - \bar{K}^0$ system. Gauge-mediated supersymmetry breaking is used to provide the needed squark mass degeneracies and A -term proportionalities; though that proves to be still insufficient for a generic Nelson-Barr model. The workable model we consider here, essentially a supersymmetric version of the aspon model, has the Nelson-Barr mass texture enforced by a $U(1)$ gauge symmetry, broken at the TeV scale. The resulting model is predictive with rich phenomenology soon to be available. Feasibility of the model considered is established by a detailed renormalization group studies.

* Talk given at PASCOS-98, Northeastern University, March 1998, and at the 4th International Workshop on Particle Phenomenology, National Sun Yat-Sen University, Taiwan June 1998 — submission for proceedings.

Gauge Mediated Supersymmetry Breaking and Spontaneous CP Violation as a Solution to the Strong CP Problem

Otto C.W. Kong

*Department of Physics and Astronomy,
University of Rochester, Rochester NY 14627-0171.*

The origin of CP violation is a major mystery, especially in relation to the strong CP problem. CP being a spontaneously broken symmetry could provide an elegant solution. However, such models have difficulty making themselves compatible with low-energy supersymmetry, which is popularly accepted as the solution to the hierarchy problem. We demonstrate that a certain class of low scale supersymmetric “Nelson-Barr” type models can solve the strong and supersymmetric CP problems while at the same time generating sufficient weak CP violation in the $K^0 - \bar{K}^0$ system. Gauge-mediated supersymmetry breaking is used to provide the needed squark mass degeneracies and A -term proportionalities; though that proves to be still insufficient for a generic Nelson-Barr model. The workable model we consider here, essentially a supersymmetric version of the aspon model, has the Nelson-Barr mass texture enforced by a U(1) gauge symmetry, broken at the TeV scale. The resulting model is predictive with rich phenomenology soon to be available. Feasibility of the model considered is established by a detailed renormalization group studies.

The sources of CP violation is still a major puzzle. While the only experimentally observed CP violating effect, in the $K^0 - \bar{K}^0$ system, is compatible with the Kabayashi-Maskawa (KM) prescription within the standard model, there are evidents for a more elaborate theory of CP violation. Among them, the strong CP problem is the most pressing. A potential strong interaction contribution to the neutron electric dipole moment and leads to the very stringent experimental constraint

$$\bar{\theta} < 10^{-9} . \quad (1)$$

Here, $\bar{\theta}$ is the physical combination of the strong CP parameter θ and the CP phase(s) of the color fermion mass matrices. In the case of the supersymmetric standard model, this is to be given by

$$\bar{\theta} = \theta - \arg \det M_q - 3 \arg M_g , \quad (2)$$

where M_q is the quark mass matrix and M_g the gluino mass. The incredible smallness of this *a priori* arbitrary parameter demands an explanation. With supersymmetry (SUSY), there are also a large number of admissible CP violating phases among the soft SUSY breaking parameters (SSBP), which are also constrained to be quite small ($\sim 10^{-3}$).

Using spontaneous CP violation to solve the strong CP problem was pioneered by Nelson and Barr¹. Extending the approach to a SUSY model could take care of the SUSY CP phases easily. Attempts to realize the Nelson-Barr mechanism in SUSY models² have, however, run up against a formidable difficulty: There generically exist potentially large 1-loop contributions to $\bar{\theta}$ in these models. As discussed at length in Ref.³, one requires an exceptionally high degree of proportionality of the soft SUSY breaking trilinear scalar couplings to their associated Yukawa couplings as well as degeneracy among the soft squark mass terms for each charge and color sector, if these contributions are to be sufficiently suppressed. The degree of proportionality and degeneracy required among the SSBP is very difficult to maintain due to the effects of renormalization.

A particularly interesting modified version of the Nelson-Barr type model is given by the aspon model⁴. Here the extra symmetry needed to enforce the Nelson-Barr mass texture is promoted to a new $U(1)$ gauge symmetry. This $U(1)_A$ gauge boson (aspon) then mediates a tree level CP violating $K^0 - \bar{K}^0$ mixing. This allows the effective KM phase to be small. A preliminary study suggested a SUSY version of the model has a much better chance to provide a feasible solution⁵. The detail model is constructed in the framework of gauge mediated SUSY breaking (GMSB), and demonstrated to work, through a careful renormalization group (RG) study⁶.

The model have CP spontaneously broken at low energies (of order a TeV). It has an extra right-handed down quark superfield $\bar{D}$ together with its mirror D coupling to the ordinary down quarks via the superpotential

$$W_d = Y_d^{ij} Q_j \bar{d}_i H_d + \mu_D D \bar{D} + \gamma^{ia} D \chi_a \bar{d}_i , \quad (3)$$

where the VEVs of the scalar components of the two χ_a 's contain a relative phase, thus breaking CP. D and χ_a have opposite $U(1)_A$ charges. The down-sector quark mass matrix can be written as

$$m_q = \begin{pmatrix} m_d & x\mu_D \mathbf{a} \\ 0 & \mu_D \end{pmatrix} , \quad (4)$$

and squark mass matrices as

$$\widetilde{M}_{RL}^2 = \begin{pmatrix} \bar{A}_d m_q (1 + \mu \tan \beta) & \bar{A}_\gamma x \mu_D \mathbf{a} \\ 0 & \bar{A}_\gamma \mu_D \end{pmatrix} + \begin{pmatrix} \langle H_d \rangle \delta A_d x \mu_D \delta A_\gamma \mathbf{c} - \gamma^{ia} \langle F_{\chi_a} \rangle & \\ 0 & \mu_D (B_D - \bar{A}_\gamma) \end{pmatrix} \quad (5)$$

$$\widetilde{M}_{RR}^2 = \tilde{m}_R^2 I + \begin{pmatrix} \delta \tilde{m}_R^2 & 0 \\ 0 & \delta \tilde{M}_R^2 \end{pmatrix} + m_q m_q^\dagger , \quad (6)$$

$$\widetilde{M}_{LL}^2 = \tilde{m}_L^2 I + \begin{pmatrix} \delta \tilde{m}_L^2 & 0 \\ 0 & \delta \tilde{M}_L^2 \end{pmatrix} + m_q^\dagger m_q , \quad (7)$$

where $a^i = \frac{1}{x\mu_D} \gamma^{ia} \langle \chi_a \rangle$, such that $\mathbf{a}^\dagger \mathbf{a} = 1$, and

$$\delta A_\gamma c^i = \frac{1}{x\mu_D} h_\gamma^{ia} \langle \chi_a \rangle - \bar{A}_\gamma a^i, \quad (8)$$

with $\mathbf{c}^\dagger \mathbf{c} = 1$. Hence, the complex phases are all contained in three-vectors $\mathbf{a}$ and $\mathbf{c}$, and the most dangerous $\langle F_{\chi_a} \rangle$'s.

The spontaneous CP violation (SCPV) is obtained from a superpotential of χ_a 's together with two oppositely charged $\bar{\chi}_a$'s and a neutral $\aleph$ superfield,

$$W_\chi = \bar{\chi}_a \mu_\chi^{ab} \chi_b + \aleph \bar{\chi}_a \lambda^{ab} \chi_b + \lambda_\aleph \aleph^3 + \mu_\aleph \aleph^2, \quad (9)$$

which generally gives a SUSY perserving vacuum with complex VEV's breaking CP. The sector is minimal for anomaly cancellation and SCPV. Without nonzero SSBP for superfields of the sector, the $\langle F_{\chi_a} \rangle$'s are then vanishing. A minimal GMSB scenario⁷ blind to the $U(1)_A$ is used to suppress proportionality violations δA_d , δA_γ , and $(B_D - \bar{A}_\gamma)$, degeneracy violations, as well as the SSBP for that lifts the zero $\langle F_{\chi_a} \rangle$'s. The latter is then estimates both analytically and diagrammatically, based on RG analysis of all SSBP. The 1-loop $\bar{\theta}$ contributions from each terms is calculated through the mass insertion method. The model predicts a measureable neutron EDM, which could be close to the present experimental bound for $x > 0.01$, with some region of the admissible x -values (of the non-SUSY version) plausibly rule out by having the $\langle F_{\chi_a} \rangle$'s too large, and also a very rich spectrum of new particles around the TeV scale.

References

1. A. Nelson, *Phys. Lett.* **136B**, 387 (1984); S. M. Barr, *ibid.* **53**, 329 (1984); *Phys. Rev.* **D 30**, 1805 (1984); A. Nelson, *Phys. Lett.* **143B**, 165 (1984).
2. S.M. Barr and A. Masiero, *Phys. Rev.* **D 38**, 366 (1988); S.M. Barr and G. Segrè, *ibid.* **D 48**, 302 (1993).
3. M. Dine, R. Leigh, and A. Kagan, *Phys. Rev.* **D 48**, 2214 (1993).
4. P.H. Frampton and T.W. Kephart, *Phys. Rev. Lett.* **66**, 1666 (1991); P.H. Frampton and D. Ng, *Phys. Rev.* **D 43**, 3034 (1991).
5. P.H. Frampton and O.C.W. Kong, *Phys. Lett.* **B402**, 297 (1997).
6. O. C. W. Kong, B. D. Wright, *Phys. Rev.* **D 58**, 015002 (1998).
7. See, for example, R. Rattazzi and U. Sarid, *Nucl. Phys.* **N501**, 297 (1997); F.M. Borzumati, WIS-96/50/Dec.-PH, hep-ph/9702307.